

**PACKAGE LEAFLET:
INFORMATION FOR THE USER**

**Lorvacs XL 1.5 mg
Prolonged-release Tablets**
Indapamide



Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Lorvacs XL 1.5 mg is and what it is used for
2. Before you take Lorvacs XL 1.5 mg
3. How to take Lorvacs XL 1.5 mg
4. Possible side effects
5. How to store Lorvacs XL 1.5 mg
6. Further information

1. WHAT LORVACS XL 1.5 MG IS AND WHAT IT IS USED FOR

This medicine is intended to reduce high blood pressure (hypertension). It is a prolonged-release tablet containing indapamide as the active ingredient. Indapamide is a diuretic. Most diuretics increase the amount of urine produced by the kidneys.

2. BEFORE YOU TAKE LORVACS XL 1.5 MG

Do not take Lorvacs XL 1.5 mg

- if you are allergic (hypersensitive) to indapamide or any other sulphonamide or to any of the other ingredients of Lorvacs XL 1.5 mg,
- if you have severe kidney disease,
- if you have severe liver disease or suffer from a condition called hepatic encephalopathy (liver problems which affect the brain and central nervous system),
- if you have low potassium levels in your blood.

Take special care with Lorvacs XL 1.5 mg

- if you have liver problems,
- if you have diabetes,
- if you suffer from gout,
- if you have any heart rhythm problems or problems with your kidneys,
- if you need to have a test to check how well your parathyroid gland is working.

You should tell your doctor if you have had photosensitivity reactions.

Your doctor may give you blood tests to check for low sodium or potassium levels or high calcium levels.

If you think any of these situations may apply to you or you have any questions or doubts about taking your medicine, you should consult your doctor or pharmacist.

Athletes should be aware that this medicine contains an active ingredient, which may give a positive reaction in doping tests.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

You should not take Lorvacs XL 1.5 mg with lithium (used to treat depression) due to the risk of increased levels of lithium in the blood.

Make sure to tell your doctor if you are taking any of the following medicines, as special care may be required:

- medicines used for heart rhythm problems (e.g. quinidine, hydroquinidine, disopyramide, amiodarone, sotalol, ibutilide, dofetilide, digitalis),
- medicines used to treat mental disorders such as depression, anxiety, schizophrenia (e.g. tricyclic antidepressants, antipsychotic drugs, neuroleptics),
- bepridil (used to treat angina pectoris, a condition causing chest pain),
- cisapride (used to treat reduced movement of the gullet and stomach),
- diphemanil (used to treat gastric problems such as ulcers, too much acid, overactive digestive system),
- sparfloxacin, moxifloxacin (antibiotics used to treat infections),
- halofantrine (antiparasitic drugs used to treat certain types of malaria),
- pentamidine (used to treat certain types of pneumonia),
- mizolastine (used to treat allergic reactions, such as hay fever),
- non-steroidal anti-inflammatory drugs for pain relief (e.g. ibuprofen) or high doses of acetylsalicylic acid,
- angiotensin converting enzyme (ACE) inhibitors (used to treat high blood pressure and heart failure),
- oral corticosteroids used to treat various conditions including severe asthma and rheumatoid arthritis,
- stimulant laxatives,
- baclofen (to treat muscle stiffness occurring in diseases such as multiple sclerosis),
- potassium-sparing diuretics (amiloride, spironolactone, triamterene),
- metformin (to treat diabetes),
- iodinated contrast media (used for tests involving X-rays),
- calcium tablets or other calcium supplements,
- ciclosporin, tacrolimus or other medicines to depress the immune system after organ transplantation, to treat autoimmune diseases, or severe rheumatic or dermatological diseases,
- tetracosactide (to treat Crohn's disease).

Pregnancy and breast-feeding

Ask your doctor or pharmacist for advice before taking any medicine.

This medicine is not recommended during pregnancy. When a pregnancy is planned or confirmed, the switch to an alternative treatment should be initiated as soon as possible. Please tell your doctor if you are pregnant or wish to become pregnant. The active ingredient is excreted in milk. Breastfeeding is not advisable if you are taking this medicine.

Driving and using machines

This medicine can cause side effects such as dizziness or tiredness due to lowering of the blood pressure (see section 4). These side effects are more likely to occur after initiation of the treatment and after dose increases. If this occurs, you should refrain from driving and other activities requiring alertness.

Important information about some of the ingredients of Lorvacs XL 1.5 mg

This medicine contains lactose monohydrate. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. HOW TO TAKE LORVACS XL 1.5 MG

Always take Lorvacs XL 1.5 mg exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

The usual dose is one tablet each day, preferably in the morning. The tablets can be taken with or without food. They should be swallowed whole



LOCATION : Indrad		COUNTRY : UK		Supersedes A/W No.:	
SIZE : 150 x 300 mm_Front Side		CODE : 8026619-9703		DATE : 25-01-2011	
REMARK :					
SUBSTRATE :					
Activities	Department	Name	Signature	Date	
Prepared By	Pkg.Dev				
Reviewed By	Pkg.Dev				
Reviewed By	RA				
Approved By	CQA				
This colour proof is not colour binding. Follow Pantone shade reference for actual colour matching.					



with water. Do not crush or chew them. Treatment for high blood pressure is usually life-long.

If you take more Lorcvas XL 1.5 mg than you should

If you have taken too many tablets, contact your doctor or pharmacist immediately.

A very large dose of Lorcvas XL 1.5 mg could cause nausea (feeling sick), vomiting, low blood pressure, cramps, dizziness, drowsiness, confusion and changes in the amount of urine produced by the kidneys.

If you forget to take Lorcvas XL 1.5 mg

If you forget to take a dose of your medicine, take the next dose at the usual time. Do not take a double dose to make up for the forgotten dose.

If you stop taking Lorcvas XL 1.5 mg

As the treatment for high blood pressure is usually life-long, you should discuss with your doctor before stopping this medicinal product.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Lorcvas XL 1.5 mg can cause side effects, although not everybody gets them.

These can include:

Commonly (less than 1 patient in 10 but more than 1 in 100):

- Low potassium in the blood, which may cause muscle weakness, allergic reactions, mainly dermatological, such as skin rashes

Uncommonly (less than 1 patient in 100 but more than 1 in 1000):

- Vomiting, purpura (red pinpoint spots on skin) in subjects with a tendency to allergic and asthmatic reactions.

Rarely (less than 1 patient in 1000 but more than 1 in 10,000):

- Feeling of tiredness, dizziness, headache, pins and needles (paresthesia);
- Nausea (feeling sick), constipation, dry mouth;
- Increased risk of dehydration in the elderly and in patients suffering from heart failure.

Very rarely (less than 1 patient in 10,000):

- Heart rhythm irregularities (causing palpitations, feeling of the heart pounding), low blood pressure;
- Kidney disease (causing symptoms of tiredness, increased need to urinate, itchy skin, feeling sick, swollen extremities);
- Pancreatitis (inflammation of the pancreas which causes upper abdominal pain), abnormal liver function (with symptoms such as tiredness, loss of appetite, feeling or being sick, swollen extremities, yellow skin). In cases of liver failure, there is a possibility of getting hepatic encephalopathy (liver problems which affect the brain and central nervous system);
- Changes in blood cells, such as thrombocytopenia (decrease in the number of platelets which causes easy bruising and nasal bleeding), leucopenia (decrease in white blood cells which may cause unexplained fever, soreness of the throat or other flu-like symptoms – if this occurs, contact your doctor) and anaemia (decrease in red blood cells);
- Angioedema and/or urticaria, severe skin manifestations. Angioedema is characterised by swelling of the skin around the eyes, lips, hands or feet. It may cause swelling of the throat, tongue or airways resulting in shortness of breath or difficulty in swallowing. If this occurs, contact your doctor immediately.

If you suffer from systemic lupus erythematosus (a disorder of the immune system leading to inflammation and damage to the joints, tendons and organs with symptoms including skin rashes, tired-

ness, loss of appetite, weight gain and joint pain), this might get worse. Cases of photosensitivity reactions (change in skin appearance) after exposure to the sun or artificial UVA have also been reported.

Some changes may occur in your blood and your doctor may need to give you blood tests to check your condition. The following changes in your blood test results may occur:

- low potassium in the blood,
- low sodium in the blood that may lead to dehydration and low blood pressure,
- increase in uric acid, a substance which may cause or worsen gout - painful joint(s) especially in the feet,
- increase in blood glucose levels in diabetic patients,
- increase of calcium in blood.

If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

5. HOW TO STORE LORCVAS XL 1.5 MG

Keep out of the reach and sight of children.

Do not use this medicine after the expiry data which is stated on the carton and blister. The expiry date refers to the last day of that month.

This product does not require any special storage conditions.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

What Lorcvas XL 1.5 mg contains

The active substance is indapamide. Each tablet contains 1.5 mg of indapamide.

The other ingredients are lactose monohydrate, pregelatinised maize starch, hypromellose, silica colloidal anhydrous, magnesium stearate, macrogol 6000, titanium dioxide (E 171).

What Lorcvas XL 1.5 mg looks like and contents of the pack

White to off-white, round shaped, biconvex, prolonged-release film-coated tablets which come in boxes of 10, 14, 15, 20, 30, 50, 60, 90 and 100. Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Torrent Pharma (UK) Ltd.

Unit 4, Charlwood Court,
County Oak Way
Crawley
West Sussex. RH11 7XA
United Kingdom

Manufacturer

Torrent Pharma (UK) Ltd.

Unit 4, Charlwood Court,
County Oak Way
Crawley
West Sussex. RH11 7XA
United Kingdom

This leaflet was last approved in 01/2011.

PL 36687/0001

8026619-9703

LOCATION : Indrad	COUNTRY : UK	Supersedes A/W No.:		
SIZE : 150 x 300 mm_Back Side	CODE : 8026619-9703	DATE : 25-01-2011		
REMARK :				
SUBSTRATE :				
Activities	Department	Name	Signature	Date
Prepared By	Pkg.Dev			
Reviewed By	Pkg.Dev			
Reviewed By	RA			
Approved By	CQA			
This colour proof is not colour binding. Follow Pantone shade reference for actual colour matching.				

